

Apple

AP<sup>®</sup> CALCULUS AB  
2008 SCORING GUIDELINES (Form B)

Question 2

For time  $t \geq 0$  hours, let  $r(t) = 120(1 - e^{-10t^2})$  represent the speed, in kilometers per hour, at which a car travels along a straight road. The number of liters of gasoline used by the car to travel  $x$  kilometers is modeled by  $g(x) = 0.05x(1 - e^{-x/2})$ .

- (a) How many kilometers does the car travel during the first 2 hours?  
 (b) Find the rate of change with respect to time of the number of liters of gasoline used by the car when  $t = 2$  hours. Indicate units of measure.  
 (c) How many liters of gasoline have been used by the car when it reaches a speed of 80 kilometers per hour?

(a)  $\int_0^2 r(t) dt = 206.370$  kilometers

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b)  $\frac{dg}{dt} = \frac{dg}{dx} \cdot \frac{dx}{dt}; \quad \frac{dx}{dt} = r(t)$   
 $\left. \frac{dg}{dt} \right|_{t=2} = \left. \frac{dg}{dx} \right|_{x=206.370} \cdot r(2)$   
 $= (0.050)(120) = 6$  liters/hour

3 :  $\begin{cases} 2 : \text{uses chain rule} \\ 1 : \text{answer with units} \end{cases}$

- (c) Let  $T$  be the time at which the car's speed reaches 80 kilometers per hour.

Then,  $r(T) = 80$  or  $T = 0.331453$  hours.

At time  $T$ , the car has gone

$x(T) = \int_0^T r(t) dt = 10.794097$  kilometers

and has consumed  $g(x(T)) = 0.537$  liters of gasoline.

4 :  $\begin{cases} 1 : \text{equation } r(t) = 80 \\ 2 : \text{distance integral} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2005 SCORING GUIDELINES (Form B)**

**Question 3**

A particle moves along the  $x$ -axis so that its velocity  $v$  at time  $t$ , for  $0 \leq t \leq 5$ , is given by

$v(t) = \ln(t^2 - 3t + 3)$ . The particle is at position  $x = 8$  at time  $t = 0$ .

- (a) Find the acceleration of the particle at time  $t = 4$ .  
 (b) Find all times  $t$  in the open interval  $0 < t < 5$  at which the particle changes direction. During which time intervals, for  $0 \leq t \leq 5$ , does the particle travel to the left?  
 (c) Find the position of the particle at time  $t = 2$ .  
 (d) Find the average speed of the particle over the interval  $0 \leq t \leq 2$ .

(a)  $a(4) = v'(4) = \frac{5}{7}$

1 : answer

(b)  $v(t) = 0$   
 $t^2 - 3t + 3 = 1$   
 $t^2 - 3t + 2 = 0$   
 $(t-2)(t-1) = 0$   
 $t = 1, 2$

3 :  $\left\{ \begin{array}{l} 1 : \text{sets } v(t) = 0 \\ 1 : \text{direction change at } t = 1, 2 \\ 1 : \text{interval with reason} \end{array} \right.$

$v(t) > 0$  for  $0 < t < 1$   
 $v(t) < 0$  for  $1 < t < 2$   
 $v(t) > 0$  for  $2 < t < 5$

The particle changes direction when  $t = 1$  and  $t = 2$ .  
 The particle travels to the left when  $1 < t < 2$ .

(c)  $s(t) = s(0) + \int_0^t \ln(u^2 - 3u + 3) du$   
 $s(2) = 8 + \int_0^2 \ln(u^2 - 3u + 3) du$   
 $= 8.368$  or  $8.369$

3 :  $\left\{ \begin{array}{l} 1 : \int_0^2 \ln(u^2 - 3u + 3) du \\ 1 : \text{handles initial condition} \\ 1 : \text{answer} \end{array} \right.$


(d)  $\frac{1}{2} \int_0^2 |v(t)| dt = 0.370$  or  $0.371$

2 :  $\left\{ \begin{array}{l} 1 : \text{integral} \\ 1 : \text{answer} \end{array} \right.$

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES (Form B)**

**Question 2**

A particle moves along the  $x$ -axis so that its velocity  $v$  at time  $t \geq 0$  is given by  $v(t) = \sin(t^2)$ . The graph of  $v$  is shown above for  $0 \leq t \leq \sqrt{5\pi}$ . The position of the particle at time  $t$  is  $x(t)$  and its position at time  $t = 0$  is  $x(0) = 5$ .



- (a) Find the acceleration of the particle at time  $t = 3$ .  
 (b) Find the total distance traveled by the particle from time  $t = 0$  to  $t = 3$ .  
 (c) Find the position of the particle at time  $t = 3$ .  
 (d) For  $0 \leq t \leq \sqrt{5\pi}$ , find the time  $t$  at which the particle is farthest to the right. Explain your answer.

(a)  $a(3) = v'(3) = 6\cos 9 = -5.466$  or  $-5.467$

(b) Distance =  $\int_0^3 |v(t)| dt = 1.702$

OR

For  $0 < t < 3$ ,  $v(t) = 0$  when  $t = \sqrt{\pi} = 1.77245$  and

$t = \sqrt{2\pi} = 2.50663$

$x(0) = 5$

$x(\sqrt{\pi}) = 5 + \int_0^{\sqrt{\pi}} v(t) dt = 5.89483$

$x(\sqrt{2\pi}) = 5 + \int_0^{\sqrt{2\pi}} v(t) dt = 5.43041$

$x(3) = 5 + \int_0^3 v(t) dt = 5.77356$

$|x(\sqrt{\pi}) - x(0)| + |x(\sqrt{2\pi}) - x(\sqrt{\pi})| + |x(3) - x(\sqrt{2\pi})| = 1.702$

(c)  $x(3) = 5 + \int_0^3 v(t) dt = 5.773$  or  $5.774$

(d) The particle's rightmost position occurs at time  $t = \sqrt{\pi} = 1.772$ .

The particle changes from moving right to moving left at those times  $t$  for which  $v(t) = 0$  with  $v(t)$  changing from positive to negative, namely at

$t = \sqrt{\pi}, \sqrt{3\pi}, \sqrt{5\pi}$  ( $t = 1.772, 3.070, 3.963$ ).

Using  $x(T) = 5 + \int_0^T v(t) dt$ , the particle's positions at the times it

changes from rightward to leftward movement are:

$T: 0 \quad \sqrt{\pi} \quad \sqrt{3\pi} \quad \sqrt{5\pi}$

$x(T): 5 \quad 5.895 \quad 5.788 \quad 5.752$

The particle is farthest to the right when  $T = \sqrt{\pi}$ .

1 :  $a(3)$

2 :  $\begin{cases} 1 : \text{setup} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 2 \begin{cases} 1 : \text{integrand} \\ 1 : \text{uses } x(0) = 5 \end{cases} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 1 : \text{sets } v(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 4**

A particle moves along the  $x$ -axis with position at time  $t$  given by  $x(t) = e^{-t} \sin t$  for  $0 \leq t \leq 2\pi$ .

- (a) Find the time  $t$  at which the particle is farthest to the left. Justify your answer.  
 (b) Find the value of the constant  $A$  for which  $x(t)$  satisfies the equation  $Ax''(t) + x'(t) + x(t) = 0$  for  $0 < t < 2\pi$ .

- (a)  $x'(t) = -e^{-t} \sin t + e^{-t} \cos t = e^{-t}(\cos t - \sin t)$   
 $x'(t) = 0$  when  $\cos t = \sin t$ . Therefore,  $x'(t) = 0$  on  
 $0 \leq t \leq 2\pi$  for  $t = \frac{\pi}{4}$  and  $t = \frac{5\pi}{4}$ .

The candidates for the absolute minimum are at

$t = 0, \frac{\pi}{4}, \frac{5\pi}{4},$  and  $2\pi$ .

$t$	$x(t)$
0	$e^0 \sin(0) = 0$
$\frac{\pi}{4}$	$e^{-\frac{\pi}{4}} \sin\left(\frac{\pi}{4}\right) > 0$
$\frac{5\pi}{4}$	$e^{-\frac{5\pi}{4}} \sin\left(\frac{5\pi}{4}\right) < 0$
$2\pi$	$e^{-2\pi} \sin(2\pi) = 0$

The particle is farthest to the left when  $t = \frac{5\pi}{4}$ .

- (b)  $x''(t) = -e^{-t}(\cos t - \sin t) + e^{-t}(-\sin t - \cos t)$   
 $= -2e^{-t} \cos t$

$$\begin{aligned} Ax''(t) + x'(t) + x(t) &= A(-2e^{-t} \cos t) + e^{-t}(\cos t - \sin t) + e^{-t} \sin t \\ &= (-2A + 1)e^{-t} \cos t \\ &= 0 \end{aligned}$$

Therefore,  $A = \frac{1}{2}$ .


$$5 : \begin{cases} 2 : x'(t) \\ 1 : \text{sets } x'(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

$$4 : \begin{cases} 2 : x''(t) \\ 1 : \text{substitutes } x''(t), x'(t), \text{ and } x(t) \\ \quad \text{into } Ax''(t) + x'(t) + x(t) \\ 1 : \text{answer} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB  
2005 SCORING GUIDELINES**

**Question 5**

A car is traveling on a straight road. For  $0 \leq t \leq 24$  seconds, the car's velocity  $v(t)$ , in meters per second, is modeled by the piecewise-linear function defined by the graph above.



- (a) Find  $\int_0^{24} v(t) dt$ . Using correct units, explain the meaning of  $\int_0^{24} v(t) dt$ .
- (b) For each of  $v'(4)$  and  $v'(20)$ , find the value or explain why it does not exist. Indicate units of measure.
- (c) Let  $a(t)$  be the car's acceleration at time  $t$ , in meters per second per second. For  $0 < t < 24$ , write a piecewise-defined function for  $a(t)$ .
- (d) Find the average rate of change of  $v$  over the interval  $8 \leq t \leq 20$ . Does the Mean Value Theorem guarantee a value of  $c$ , for  $8 < c < 20$ , such that  $v'(c)$  is equal to this average rate of change? Why or why not?

- (a)  $\int_0^{24} v(t) dt = \frac{1}{2}(4)(20) + (12)(20) + \frac{1}{2}(8)(20) = 360$   
The car travels 360 meters in these 24 seconds.

2 :  $\begin{cases} 1 : \text{value} \\ 1 : \text{meaning with units} \end{cases}$

- (b)  $v'(4)$  does not exist because
- $$\lim_{t \rightarrow 4^-} \left( \frac{v(t) - v(4)}{t - 4} \right) = 5 \neq 0 = \lim_{t \rightarrow 4^+} \left( \frac{v(t) - v(4)}{t - 4} \right).$$
- $$v'(20) = \frac{20 - 0}{16 - 24} = -\frac{5}{2} \text{ m/sec}^2$$

3 :  $\begin{cases} 1 : v'(4) \text{ does not exist, with explanation} \\ 1 : v'(20) \\ 1 : \text{units} \end{cases}$

- (c) 
$$a(t) = \begin{cases} 5 & \text{if } 0 < t < 4 \\ 0 & \text{if } 4 < t < 16 \\ -\frac{5}{2} & \text{if } 16 < t < 24 \end{cases}$$
  
 $a(t)$  does not exist at  $t = 4$  and  $t = 16$ .

2 :  $\begin{cases} 1 : \text{finds the values } 5, 0, -\frac{5}{2} \\ 1 : \text{identifies constants with correct intervals} \end{cases}$


- (d) The average rate of change of  $v$  on  $[8, 20]$  is
- $$\frac{v(20) - v(8)}{20 - 8} = -\frac{5}{6} \text{ m/sec}^2.$$
- No, the Mean Value Theorem does not apply to  $v$  on  $[8, 20]$  because  $v$  is not differentiable at  $t = 16$ .

2 :  $\begin{cases} 1 : \text{average rate of change of } v \text{ on } [8, 20] \\ 1 : \text{answer with explanation} \end{cases}$

AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES (Form B)

Question 4

The rate, in calories per minute, at which a person using an exercise machine burns calories is modeled by the function  $f$ . In the figure above,  $f(t) = -\frac{1}{4}t^3 + \frac{3}{2}t^2 + 1$  for  $0 \leq t \leq 4$  and  $f$  is piecewise linear for  $4 \leq t \leq 24$ .



- (a) Find  $f'(22)$ . Indicate units of measure.
- (b) For the time interval  $0 \leq t \leq 24$ , at what time  $t$  is  $f$  increasing at its greatest rate? Show the reasoning that supports your answer.
- (c) Find the total number of calories burned over the time interval  $6 \leq t \leq 18$  minutes.
- (d) The setting on the machine is now changed so that the person burns  $f(t) + c$  calories per minute. For this setting, find  $c$  so that an average of 15 calories per minute is burned during the time interval  $6 \leq t \leq 18$ .

(a)  $f'(22) = \frac{15 - 3}{20 - 24} = -3$  calories/min/min

(b)  $f$  is increasing on  $[0, 4]$  and on  $[12, 16]$ .

On  $(12, 16)$ ,  $f'(t) = \frac{15 - 9}{16 - 12} = \frac{3}{2}$  since  $f$  has constant slope on this interval.

On  $(0, 4)$ ,  $f'(t) = -\frac{3}{4}t^2 + 3t$  and

$f''(t) = -\frac{3}{2}t + 3 = 0$  when  $t = 2$ . This is where  $f'$  has a maximum on  $[0, 4]$  since  $f'' > 0$  on  $(0, 2)$  and  $f'' < 0$  on  $(2, 4)$ .

On  $[0, 24]$ ,  $f$  is increasing at its greatest rate when  $t = 2$  because  $f'(2) = 3 > \frac{3}{2}$ .

(c)  $\int_6^{18} f(t) dt = 6(9) + \frac{1}{2}(4)(9 + 15) + 2(15)$   
 $= 132$  calories

(d) We want  $\frac{1}{12} \int_6^{18} (f(t) + c) dt = 15$ .

This means  $132 + 12c = 15(12)$ . So,  $c = 4$ .

OR

Currently, the average is  $\frac{132}{12} = 11$  calories/min.

Adding  $c$  to  $f(t)$  will shift the average by  $c$ .

So  $c = 4$  to get an average of 15 calories/min.

1 :  $f'(22)$  and units

4 :  $\begin{cases} 1 : f' \text{ on } (0, 4) \\ 1 : \text{shows } f' \text{ has a max at } t = 2 \text{ on } (0, 4) \\ 1 : \text{shows for } 12 < t < 16, f'(t) < f'(2) \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{method} \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{setup} \\ 1 : \text{value of } c \end{cases}$

