

AP Calculus Review & Applications Kiwi Set

2007 B No calculator

6. Let f be a twice-differentiable function such that $f(2) = 5$ and $f(5) = 2$. Let g be the function given by $g(x) = f(f(x))$.
- Explain why there must be a value c for $2 < c < 5$ such that $f'(c) = -1$.
 - Show that $g'(2) = g'(5)$. Use this result to explain why there must be a value k for $2 < k < 5$ such that $g''(k) = 0$.
 - Show that if $f''(x) = 0$ for all x , then the graph of g does not have a point of inflection.
 - Let $h(x) = f(x) - x$. Explain why there must be a value r for $2 < r < 5$ such that $h(r) = 0$.

2007 A
Yes Calc

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	9	2	3	1
3	10	-4	4	2
4	-1	3	6	7

3. The functions f and g are differentiable for all real numbers, and g is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of x . The function h is given by $h(x) = f(g(x)) - 6$.
- Explain why there must be a value r for $1 < r < 3$ such that $h(r) = -5$.
 - Explain why there must be a value c for $1 < c < 3$ such that $h'(c) = -5$.
 - Let w be the function given by $w(x) = \int_1^{g(x)} f(t) dt$. Find the value of $w'(3)$.
 - If g^{-1} is the inverse function of g , write an equation for the line tangent to the graph of $y = g^{-1}(x)$ at $x = 2$.


2007 A No calculator

6. Let f be the function defined by $f(x) = k\sqrt{x} - \ln x$ for $x > 0$, where k is a positive constant.
- Find $f'(x)$ and $f''(x)$.
 - For what value of the constant k does f have a critical point at $x = 1$? For this value of k , determine whether f has a relative minimum, relative maximum, or neither at $x = 1$. Justify your answer.
 - For a certain value of the constant k , the graph of f has a point of inflection on the x -axis. Find this value of k .

2006 Form A NO Calc.

6. The twice-differentiable function f is defined for all real numbers and satisfies the following conditions:
- $$f(0) = 2, f'(0) = -4, \text{ and } f''(0) = 3.$$
- The function g is given by $g(x) = e^{ax} + f(x)$ for all real numbers, where a is a constant. Find $g'(0)$ and $g''(0)$ in terms of a . Show the work that leads to your answers.
 - The function h is given by $h(x) = \cos(kx)f(x)$ for all real numbers, where k is a constant. Find $h'(x)$ and write an equation for the line tangent to the graph of h at $x = 0$.

2006 B
Yes
Calc




3. The figure above is the graph of a function of x , which models the height of a skateboard ramp. The function meets the following requirements.
- At $x = 0$, the value of the function is 0, and the slope of the graph of the function is 0.
 - At $x = 4$, the value of the function is 1, and the slope of the graph of the function is 1.
 - Between $x = 0$ and $x = 4$, the function is increasing.
- Let $f(x) = ax^2$, where a is a nonzero constant. Show that it is not possible to find a value for a so that f meets requirement (ii) above.
 - Let $g(x) = cx^3 - \frac{x^2}{16}$, where c is a nonzero constant. Find the value of c so that g meets requirement (ii) above. Show the work that leads to your answer.
 - Using the function g and your value of c from part (b), show that g does not meet requirement (iii) above.
 - Let $h(x) = \frac{x^n}{k}$, where k is a nonzero constant and n is a positive integer. Find the values of k and n so that h meets requirement (ii) above. Show that h also meets requirements (i) and (iii) above.

NO
CALC

x	0	$0 < x < 1$	1	$1 < x < 2$	2	$2 < x < 3$	3	$3 < x < 4$
$f(x)$	-1	Negative	0	Positive	2	Positive	0	Negative
$f'(x)$	4	Positive	0	Positive	DNE	Negative	-3	Negative
$f''(x)$	-2	Negative	0	Positive	DNE	Negative	0	Positive

2005 A

4. Let f be a function that is continuous on the interval $[0, 4]$. The function f is twice differentiable except at $x = 2$. The function f and its derivatives have the properties indicated in the table above, where DNE indicates that the derivatives of f do not exist at $x = 2$.
- For $0 < x < 4$, find all values of x at which f has a relative extremum. Determine whether f has a relative maximum or a relative minimum at each of these values. Justify your answer.
 - On the axes provided, sketch the graph of a function that has all the characteristics of f .
(Note: Use the axes provided in the pink test booklet.)



- Let g be the function defined by $g(x) = \int_1^x f(t) dt$ on the open interval $(0, 4)$. For $0 < x < 4$, find all values of x at which g has a relative extremum. Determine whether g has a relative maximum or a relative minimum at each of these values. Justify your answer.
- For the function g defined in part (c), find all values of x , for $0 < x < 4$, at which the graph of g has a point of inflection. Justify your answer.